



A STUDY OF COASTAL ART TOWARDS SUSTAINABLE TOURISM IN OF GOA

Thejaswini R*, Dr. Vibha Lodhi

* Research Scholar, Dept. of Fashion Designing, Himalayan Garhwal University, Uttarakhand
Associate Professor, Dept. of Fashion Designing, Himalayan Garhwal University, Uttarakhand

ABSTRACT

Goa is a coastal paradise blessed with a long coastline of 105km in length consisting of number of pristine beaches. It is this lengthy coastline which has put Goa on the world map as an important tourist destination. To sustain long time tourism in the State there is a need of proper implementation of plans and action with regards to CRZ. The changing CRZ will not only take away the area of traditional fisher folk and bifurcate coastal zones along rural area based on population density but Goa will be lost to development making coastline a concrete jungle. The land-ocean interaction in the coastal zone produces a complex range of landforms and organisms which evolve special forms of adaptation. The 2018 notification of the MoEF which has proposed to reduce the existing HTL from 200mts to 50mts will change not only the fragile coastal zone of Goa but will put Tourism at stake. The pristine beaches will be lost for development leading to coastal hazards. A question arises whether a small State like Goa will be able to withstand the pressures of handling CRZ regulations due to influx of tourist which is far more than the total population of Goa.

Keywords:

Coastal Regulation Zone (CRZ), Coastal Zone, High Tide Line (HTL), Sustainable Tourism



INTRODUCTION

The coastal zone is particularly important from a human perspective. A large proportion of the world population is concentrated in the coastal zone, including almost all of the major cities. The State attracts tourists within and internationally due to its pristine beaches and places of worship. Tourism in Goa is generally focused in the coastal areas rather than as a hinterland activity. The sea front is marked by a combination of beaches, rocky shores and headlands. Out of 105km long coast, more than 70km comprises of linear and wide sandy beaches all backed by 1 to 10 meters high dunes; sandy pockets and secluded coves backed by rocky cliff are also found (Mascarenhas, 2005). Till early 70's, coastline remained pristine. Human pressure on the coastal ecosystem started in late 1970's when tourism became a major source of revenue. Thus, leading to a substantial change in the coastal area. Unchecked tourism pressure resulted in 80% of the urban growth in coastal talukas of Goa. The main purpose CRZ notification was to control and minimize environmental damage to coastal ecosystem. Ministry of Environment and Forest, Government of India enacted the Coastal Regulation Zone (CRZ) notification issued under Environment Protection Act of 1986, which was notified through Gazette in 1991. The No development zone was confined up to 500mts from HTL in the 1991, with respect to imposed restrictions on setting up and expansion of Industries, operations or processes etc., in the said Coastal Regulation Zone (CRZ) (Government of India, 2018). Subsequently, the CRZ was brought to 200mts from HTL through a notification in 2011. Recently, the new affirming rule of reducing HTL from 200mts to 50mts will procure the ecological area into urbanization. Monitoring the LULC change along the Coastal Regulation Zone (CRZ) areas is essential to understand the current status of ecologically sensitive areas in order to conserve the ecosystems from deleterious impacts (Guru B., 2014). In recent era, Remote sensing and GIS techniques are excessively used to inventory, monitor and management of natural resources in the coastal areas due to its repetitive, multispectral and synoptic nature. Geographic Information Systems (GIS) at other side provide suitable platform for data analysis, update and retrieval (Dale, 1997). Space borne remotely sensed data may be particularly useful in developing countries where recent and reliable spatial information is lacking (Deng et al., 2009). These tools will help in studying the change in any form of land use through both the external forces and the pressure built-up within the system.



Database and Methodology: The present study is a collaboration of primary and secondary data. Primary data consist of opinions gathered, personal interviews, visits to North Goa Coastline and tourist spots. Various research documents, research institutes, satellite images, multi-temporal satellite data has been used as a secondary source (Table 1). The required satellite imageries were obtained from <http://earthexplorer.usgs.gov>. Level 1 classification has been followed to determine the coast as per the guidelines of Ministry of Environment, Forest and Climate Change and National Centre for Sustainable Coastal Management (NCSCM). ERDAS Imagine 14 version, a strong GIS desktop application having powerful potentials in image processing have been used for classifying land use land cover classes and comparative study of patio-temporal changes of an area. Google earth has been used as a base reference to do the classification. HTL data of 2003 has been opted from National Institute of Oceanography, Dona Paula, Goa. Toposheets also has been used as a base map to validate coastal landscape. Processing satellite utilized with geometric calculation and symmetrical difference field option to understand land cover units. The data is presented in terms of GIS based maps, tables and graphs.

Table 1: Data Source

Sr. No.	Satellite	Sensor	Acquisition Date	Resolution	Data Source
1	LANDSAT 4-5	TM	17-02-1991	30m	USGS, Earth Explorer
2	LANDSAT08	OLI TIRS	19-02-2015	30m	

Result and Discussion

Coastal Zone is ecologically sensitive and it is a confluence of land and water. Coastal Zone Management is a major consternation for decision-makers, land use planners and environmentalists. Coastal zones fluctuate between the land-water interfaces, due to influences of climate, physiography, ecology on land and water. All the tourism related activities in the coastal areas have given rise to violations of CRZ. Implementation of CRZ regulations on the part of the government will lead to loss of fragile coastal environment which is the major attraction of the State. The land use land cover change along the coastline is an peer view to appreciate the spatial distribution and CRZ violation from HTL.

Land use Land Cover Change with A Buffer From HTL: 1991 and 2015

The tabulated data, figures and maps were generated and used for further analysis to understand the changing scenario of LULC and CRZ violation along the coast for the period of 1991 and 2015.

Table 2: LULC Change Detection: CRZ Violation Within 200mts and 50mts from HTL of Pernem

Pernem (Area in sq.km.)								
Years	Classes/ CRZ zones	Built-up	Agriculture	Vegetation Cover	Water Bodies	Others	Category- wise Total	Overall Total
1991	200mts from HTL	0.63	0.069	0.864	0.239	0.994	2.797	5.59
	50mts from HTL	0.15	0	0.181	0.051	0.330	0.710	
	50mts- 200mts	0.48	0.069	0.683	0.187	0.665	2.087	
2015	200mts from HTL	1.29	0.01	0.92	0.14	0.44	2.795	5.59
	50mts from HTL	0.28	0	0.18	0.03	0.23	0.710	
	50mts- 200mts	1.01	0.01	0.75	0.11	0.21	2.08	




Figure 1: Graph, Pernem Coastal Area in % Figure 2: Graph, Pernem Coastal Area in %




Figure 3: LULC with CRZ violation of Bardez for 1991 and 2015

It can be inferred (table 2 and figure 2, 3 and 4) that built up and vegetation has increased over a period of 1991 to 2015 from 23% to 45 % and 30 % to 33% respectively. The decline in agriculture and other category can be observed from the LULC. The implication of No development zone of 50mts from HTL shows that it will procure loss to almost 32 percent of the vegetation. As the remaining area from 50mts to 200mts will be probably converted to developmental zone. Similarly, the already existence of pollution near coast will increase in abundance. The minor existence of agricultural land turning into commercial zone has highest implications.

Table 3: LULC Change Detection: CRZ Violation Within 200mts and 50mts from HTL of Bardez.

Bardez (Area in sq.km.)								
Years	Classes/ CRZ zones	Built-up	Agriculture	Vegetation Cover	Water Bodies	Others	Total	Overall Total
1991	200mts from HTL	0.83	0.04	1.55	0.19	5.21	7.82	15.64
	50mts from HTL	0.16	0.00	0.15	0.14	1.48	1.93	
	50mts- 200mts	0.67	0.04	1.40	0.05	3.73	5.89	
2015	200mts from HTL	2.46	0.01	1.41	0.18	3.77	7.82	15.64
	50mts from HTL	0.40	0.00	0.17	0.13	1.23	1.93	
	50mts- 200mts	2.06	0.01	1.24	0.05	2.54	5.89	




Figure 4: Graph, Bardez Coastal Area in % Figure 5: Graph, Bardez Coastal Area in %




Figure 6: LULC with CRZ violation of Bardez for 1991 and 2015



The violation of CRZ can be clearly seen (figure 7), even though there was No development zone under HTL from 200mts for Bardez. In spite of decreasing the commercial or residential developmental activity, the built-up has increased from 11% to 31% respectively from the year 1991 to 2015. This includes permanent residents and temporary shacks. Whereas, rest of the classified areas shows decline in the percentage of spatial cover as per the 200mts buffer. The 50mts from HTL shows that built-up and vegetation 12% and 1% respectively. The study shows that if the CRZ declines from 200mts to 50mts the presence of vegetation and other category beyond 50mts of around 21% while 43% can be susceptible to be used for commercialization (table 3, figure 5 and 6).

CONCLUSION

Man and Environment interaction are more interactive in the coastal ecosystems and its effects are seen on basically on coastline. The study of coastline of Pernem and Bardez taluka has a grave violation not only in terms of geomorphology but also with reference to CRZ. The recent notification has changed this scenario completely. The long considered no development zone of the coastal areas will be brought under development bring the public spaces into private domain. The general public, be it locals or tourists will not be able to get access to the beaches. Therefore, the question arises as to whether the Government has any right to stop people from enjoying and appreciating the beauty of the nature. In an effort to attract more and more tourist and to earn more profit sprawling resort are built by bulldozing thousands of Casuarina trees along the coast beaches. These resorts have destroyed scenic beauty of a place and ecology. The study clearly showed the loss of natural habitat at the higher scale. The 35 percent of vegetation in Pernem and 21 percent of vegetation in Bardez are more susceptible to losses of harsh commercialization. Similarly, Commercialization and residential needs of the beaches will not only led to erosion but also CRZ violation which in long run could affect beach formation itself.



REFERENCES

- 1) Dale, V. H., 2018. The relationship between land-use changes and climate change. [Online][http://www.esajournals.org/doi/abs/10.1890/10510761\(1997\)007%5B0753%3ATRBLUC%5D2.0.CO%3B2](http://www.esajournals.org/doi/abs/10.1890/10510761(1997)007%5B0753%3ATRBLUC%5D2.0.CO%3B2) [Accessed 19 August 2014].
- 2) Deng, J. S., Wang, K., Hong, Y. & Qi, J. G., 2009. Spatio-temporal dynamics and evolution of land use change and landscape pattern in response to rapid urbanization. [Online] <http://www.sciencedirect.com/science/article/pii/S0169204609000814> [Accessed 21 August 2014].
- 3) Guru B., 2014. Study the Land use and Land cover changes and CRZ in the coastal area of Ganjam district, Odisha. [Online] <https://www.researchgate.net/publication/270277077> [Accessed 12 September 2017]
- 4) Mascarenhas A., 2015. The coastal Regulation Zone of Goa: Oceanographic, Environmental and Social perspectives. [Online] <http://www.ias.ac.in/dec251999/articles16.htm> [Accessed 12 September 2015]
- 5) Reports Government of India, (2018): Draft CRZ notification., Government of India Ministry of Environment, Forest and Climate Change. [Online] <https://www.goa.gov.in> [Accessed 12 September 2018]
- 6) National Institute of Oceanography, 2013. Demarcation of High Tide Line, Low Tide Line and CRZ boundaries in the Coastal Regulation Zone of the state of Goa (North). National Institute of Oceanography (NIO), Dona Paula Goa.